

Attorney Docket No.: DEX-0075  
Inventors: Macina and Sun  
Serial No.: 09/618,596  
Filing Date: July 17, 2000  
Page 4

#### REMARKS

Claims 1-5 are pending in the instant application. Claims 1-5 have been rejected under 35 U.S.C. § 112, first paragraph, as failing to comply with the enablement requirement. The Examiner suggests that the specification does not provide evidence supporting the effectiveness of using any CSG in the claimed methods. More specifically, the Examiner suggests that the specification does not set forth enabling disclosure supporting the occurrence of SEQ ID NO:1/Cln106 in colon cancer. Further, the Examiner suggests that the specification does not support the use of SEQ ID NO:1 in methods of diagnosing metastases, staging and monitoring of colon cancer.

Applicants respectfully traverse this rejection.

MPEP § 2164.04 is quite clear; a specification disclosure which contains a teaching of the manner and process of making and using an invention in terms which correspond in scope to those used in describing and defining the subject matter sought to be patented must be taken in compliance with the enablement requirement of 35 U.S.C. § 112, first paragraph, unless there is reason to doubt the objective truth of statements contained therein which must be relied on for enabling support. Detailed teachings for detecting/diagnosing the presence of cancer are set forth in the

Attorney Docket No.: DEX-0075  
Inventors: Macina and Sun  
Serial No.: 09/618,596  
Filing Date: July 17, 2000  
Page 5

instant application at page 7, line 27, through page 9, line 11. Detailed teachings for staging of colon cancer via the present invention are described in the specification at page 9, lines 12-25. Detailed teachings for monitoring for the onset of metastasis and/or monitoring for changes in the staging of colon cancer via the present invention are also described in the specification at page 9, line 26, through page 10, line 22. Further, assay techniques for detecting levels of CSGs for use in the methods of the present invention are described in detail in the specification at page 11, line 20, through page 14, line 30. Accordingly, the instant specification clearly contains a teaching of the manner and process of using the invention in terms which correspond in scope to those used in describing and defining the subject matter sought to be patented.

Further, in accordance with MPEP § 2164.04, in order to make a lack of enablement rejection, the Examiner has the initial burden to establish a reasonable basis to question the enablement provided for the claimed invention. As stated by the court in *In re Marzocchi*, 439 F.2d 220, 169 USPQ 367 (CCPA 1971), "[i]t is incumbent upon the Patent Office, whenever a rejection on this basis is made, to explain why it doubts the truth or accuracy of any statement in a supporting disclosure and to back up assertions

Attorney Docket No.: DEX-0075  
Inventors: Macina and Sun  
Serial No.: 09/618,596  
Filing Date: July 17, 2000  
Page 6

of its own with acceptable evidence or reasoning which is inconsistent with the contested statement." Reasoning provided by the Examiner in the instant rejection is insufficient to meet this burden as none of the evidence is directed to detection of the CSG Cln106.

However, in an earnest effort to advance the prosecution of this case Applicants have canceled claims 2 through 5. Applicants have also amended claim 1 to be drawn to a method for detecting the presence of cancer. Support for this amendment is provided in the specification at page 3, line 15 and page 6, lines 29-34. Thus, no new matter is added by these amendments.

Further, with respect to claim 1, Applicants respectfully point out that the instant application is a continuation-in-part of PCT/US99/10498. A courtesy copy of this PCT Application is provided herewith. As evidenced by data presented in this PCT application at pages 16-20, Cln106 (disclosed therein as SEQ ID NO:3) is overexpressed in 65% of the colon matching samples. Thus, as demonstrated by the parent patent application of the instant continuation-in-part application, the CSG Cln106 is a diagnostic marker for colon cancer.

This evidence establishes that there is no reason to doubt the objective truth of statements in the instant specification

Attorney Docket No.: DEX-0075  
Inventors: Macina and Sun  
Serial No.: 09/618,596  
Filing Date: July 17, 2000  
Page 7

regarding use of CSG Cln106 as a diagnostic marker for colon cancer.


Thus, the instant specification, which provides detailed teachings of methods for making and using the instant claimed invention meets the enablement requirements of 35 U.S.C. § 112, first paragraph.

Withdrawal of this rejection under 35 U.S.C. § 112, first paragraph is respectfully requested in light of the amendments to the claims and the above remarks.

#### Conclusion

Applicants believe that the foregoing comprises a full and complete response to the Office Action of record. Accordingly, favorable reconsideration and subsequent allowance of the pending claims is earnestly solicited.

Respectfully submitted,



Kathleen A. Tyrrell  
Registration No. 38,350

Date: February 2, 2004

Licata & Tyrrell P.C.  
66 E. Main Street  
Marlton, New Jersey 08053  
(856) 810-1515

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : <b>C12Q 1/68, G01N 33/53, 33/574</b>	<b>A1</b>	(11) International Publication Number: <b>WO 99/60161</b> (43) International Publication Date: 25 November 1999 (25.11.99)
(21) International Application Number: <b>PCT/US99/10498</b> (22) International Filing Date: <b>12 May 1999 (12.05.99)</b> (30) Priority Data: <b>60/086,266 21 May 1998 (21.05.98) US</b> (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application <b>US 60/086,266 (CIP)</b> <b>Filed on 21 May 1998 (21.05.98)</b> (71) Applicant (for all designated States except US): <b>DIADEXUS LLC [US/US]; 3303 Octavius Drive, Santa Clara, CA 95054 (US).</b> (72) Inventors; and (75) Inventors/Applicants (for US only): <b>MACINA, Roberto, A. [AR/US]; 4118 Crescendo Avenue, San Jose, CA 95136 (US). YANG, Fei [CN/US]; Apartment 204, 18375 Caminito Cantilena, San Diego, CA 92128 (US). SUN, Yongming [CN/US]; Apartment 260, 869 S. Winchester Boulevard, San Jose, CA 92128 (US).</b>	(74) Agents: <b>LICATA, Jane, Massey et al.; Law Offices of Jane Massey Licata, 66 E. Main Street, Marlton, NJ 08053 (US).</b> (81) Designated States: <b>CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</b>  <b>Published</b> <i>With international search report.</i>	
(54) Title: <b>A NOVEL METHOD OF DIAGNOSING, MONITORING, AND STAGING COLON CANCER</b> (57) Abstract  <b>The present invention provides a new method for detecting, diagnosing, monitoring, staging, and prognosticating colon cancer.</b>		

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BV	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WO 99/60161

PCT/US99/10498

- 1 -

A NOVEL METHOD OF DIAGNOSING,  
MONITORING, AND STAGING COLON CANCER

FIELD OF THE INVENTION

This invention relates, in part, to newly developed assays for detecting, diagnosing, monitoring, staging, and prognosticating cancers, particularly colon cancer.

BACKGROUND OF THE INVENTION

Colon cancer is the second most frequently diagnosed malignancy in the United States. Cancer of the gastrointestinal tract, especially colon cancer, is a highly treatable and often a curable disease when localized to the bowel. However, currently colon cancer is the second most common cause of cancer death. Surgery is the primary treatment and results in cure in approximately 50% of patients. Recurrence following surgery is a major problem and often is the ultimate cause of death. The prognosis of colon cancer is clearly related to the degree of penetration of the tumor through the bowel wall and the presence or absence of nodal involvement. These two characteristics form the basis for all staging systems developed for this disease. Bowel obstruction and bowel perforation are indicators of poor prognosis. Elevated pretreatment serum levels of carcinoembryonic antigen (CEA) and carbohydrate antigen 19-9 (CA 19-9) also have negative prognostic significance.

Because of the frequency of the disease (approximately 160,000 new cases of colon cancer per year), the identification of high-risk groups, the demonstrated slow growth of primary lesions, the better survival of early-stage lesions, and the relative simplicity and accuracy of screening tests, screening for colon cancer should be a part of routine

WO 99/60161

PCT/US99/10498

- 2 -

care for all adults starting at age 50, especially those with first-degree relatives with colorectal cancer.

Procedures used for detecting, diagnosing, monitoring, staging, and prognosticating colon cancer are of critical importance to the outcome of the patient. For example, patients diagnosed with early colon cancer generally have a much greater five-year survival rate as compared to the survival rate for patients diagnosed with distant metastasized colon cancer. Treatment decisions are usually made in reference to the older Dukes or the Modified Astler-Coller (MAC) classification schema for staging. However, new diagnostic methods which are more sensitive and specific for detecting early colon cancer are clearly needed.

Further, colon cancer patients must be closely monitored following initial therapy and during adjuvant therapy to determine response to therapy and to detect persistent or recurrent disease or metastasis. Thus, there is clearly a need for a colon cancer marker which is more sensitive and specific in detecting colon cancer recurrence.

Another important step in managing colon cancer is to determine the stage of the patient's disease. Stage determination has potential prognostic value and provides criteria for designing optimal therapy. Currently, pathological staging of colon cancer is preferable over clinical staging as pathological staging provides a more accurate prognosis. However, clinical staging would be preferred were the method of clinical staging at least as accurate as pathological staging because it does not depend on an invasive procedure to obtain tissue for pathological evaluation. Staging of colon cancer would be improved by detecting new markers in cells, tissues, or bodily fluids which could differentiate between different stages of invasion.

In the present invention, methods are provided for detecting, diagnosing, monitoring, staging, and



WO 99/60161

PCT/US99/10498

- 3 -

prognosticating colon cancers, particularly colon, stomach, and small intestine cancer, via nine (9) Colon Specific Genes (CSGs). The nine CSGs refer, among other things, to native proteins expressed by the genes comprising the polynucleotide sequences of any of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8 or 9. In the alternative, what is meant by the nine CSGs as used herein, means the native mRNAs encoded by the genes comprising any of the polynucleotide sequences of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8 or 9 or levels of the genes comprising any of the polynucleotide sequences of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8 or 9.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

#### SUMMARY OF THE INVENTION

Toward these ends, and others, it is an object of the present invention to provide a method for diagnosing the presence of colon cancer in a patient which comprises measuring levels of CSG in a sample of cells, tissue or bodily fluid from the patient and comparing the measured levels of CSG with levels of CSG in preferably the same cells, tissue, or bodily fluid type of a control, wherein an increase in the measured CSG levels in the patient versus levels of CSG in the control is associated with colon cancer.

Another object of the present invention is to provide a method of diagnosing metastatic colon cancer in a patient which comprises measuring CSG levels in a sample of cells,

WO 99/60161

PCT/US99/10498

- 4 -

tissue, or bodily fluid from the patient and comparing the measured CSG levels with levels of CSG in preferably the same cells, tissue, or bodily fluid type of a control, wherein an increase in measured CSG levels in the patient versus levels of CSG in the control is associated with a cancer which has metastasized.

Another object of the present invention is to provide a method of staging colon cancer in a patient which comprises identifying a patient having colon cancer, measuring levels of CSG in a sample of cells, tissues, or bodily fluid obtained from the patient, and comparing the measured CSG levels with levels of CSG in preferably the same cells, tissue or bodily fluid type of a control. An increase in measured CSG levels in the patient versus CSG levels in the control can be associated with a cancer which is progressing while a decrease or equivalent level of CSG measured in the patient versus the control can be associated with a cancer which is regressing or in remission.

Another object of the present invention is to provide a method of monitoring colon cancer in a patient for the onset of metastasis. The method comprises identifying a patient having colon cancer that is not known to have metastasized, periodically measuring levels of CSG in a sample of cells, tissues, or bodily fluid obtained from the patient, and comparing the measured CSG levels with levels of CSG in preferably the same cells, tissue, or bodily fluid type of a control, wherein an increase in measured CSG levels versus control CSG levels is associated with a cancer which has metastasized.

Yet another object of the present invention is to provide a method of monitoring the change in stage of colon cancer in a patient which comprises identifying a patient having colon cancer, periodically measuring levels of CSG in a sample of cells, tissue, or bodily fluid obtained from the patient, and comparing the measured CSG levels with levels of

WO 99/60161

PCT/US99/10498

- 5 -

CSG in preferably the same cells, tissues, or bodily fluid type of a control wherein an increase in measured CSG levels versus the control CSG levels is associated with a cancer which is progressing and a decrease in the measured CSG levels versus the control CSG levels is associated with a cancer which is regressing or in remission.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

#### DESCRIPTION OF THE INVENTION

The present invention relates to diagnostic assays and methods, both quantitative and qualitative for detecting, diagnosing, monitoring, staging, and prognosticating cancers by comparing levels of CSG with those of CSG in a normal human control. What is meant by "levels of CSG" as used herein, means levels of the native protein expressed by the genes comprising the polynucleotide sequence of any of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8 or 9. In the alternative, what is meant by "levels of CSG" as used herein, means levels of the native mRNA encoded by any of the genes comprising any of the polynucleotide sequences of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8 or 9 or levels of the gene comprising any of the polynucleotide sequence of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8 or 9. Such levels are preferably measured in at least one of, cells, tissues and/or bodily fluids, including determination of normal and abnormal levels. Thus, for instance, a diagnostic assay in accordance with the invention for

WO 99/60161

PCT/US99/10498

- 6 -

diagnosing over-expression of any one of the CSG proteins compared to normal control bodily fluids, cells, or tissue samples may be used to diagnose the presence of cancers, including colon cancer. Any of the nine CSGs may be measured alone in the methods of the invention, or all together or any combination of the nine.

By "control" it is meant a human patient without cancer and/or non cancerous samples from the patient, also referred to herein as a normal human control; in the methods for diagnosing or monitoring for metastasis, control may also include samples from a human patient that is determined by reliable methods to have colon cancer which has not metastasized.

All the methods of the present invention may optionally include measuring the levels of other cancer markers as well as CSG. Other cancer markers, in addition to CSG, useful in the present invention will depend on the cancer being tested and are known to those of skill in the art.

#### **Diagnostic Assays**

The present invention provides methods for diagnosing the presence of colon cancer by analyzing for changes in levels of CSG in cells, tissues or bodily fluids compared with levels of CSG in cells, tissues or bodily fluids of preferably the same type from a normal human control, wherein an increase in levels of CSG in the patient versus the normal human control is associated with the presence of colon cancer. Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the patient being tested has cancer is one in which cells, tissues, or bodily fluid levels of the cancer marker, such as CSG, are at least two times higher, and most preferably are at least five times higher, than in preferably the same cells, tissues, or bodily fluid of a normal human control.

The present invention also provides a method of diagnosing metastatic colon cancer in a patient having colon

WO 99/60161

PCT/US99/10498

- 7 -

cancer which has not yet metastasized for the onset of metastasis. In the method of the present invention, a human cancer patient suspected of having colon cancer which may have metastasized (but which was not previously known to have metastasized) is identified. This is accomplished by a variety of means known to those of skill in the art. For example, in the case of colon cancer, patients are typically diagnosed with colon cancer following traditional detection methods.

In the present invention, determining the presence of CSG level in cells, tissues, or bodily fluid, is particularly useful for discriminating between colon cancer which has not metastasized and colon cancer which has metastasized. Existing techniques have difficulty discriminating between colon cancer which has metastasized and colon cancer which has not metastasized and proper treatment selection is often dependent upon such knowledge.

In the present invention, the cancer marker levels measured in such cells, tissues, or bodily fluid is CSG, and are compared with levels of CSG in preferably the same cells, tissue, or bodily fluid type of a normal human control. That is, if the cancer marker being observed is just CSG in serum, this level is preferably compared with the level of CSG in serum of a normal human patient. An increase in the CSG in the patient versus the normal human control is associated with colon cancer which has metastasized.

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the cancer in the patient being tested or monitored has metastasized is one in which cells, tissues, or bodily fluid levels of the cancer marker, such as CSG, are at least two times higher, and most preferable are at least five times higher, than in preferably the same cells, tissues, or bodily fluid of a normal patient.

WO 99/60161

PCT/US99/10498

- 8 -

Normal human control as used herein includes a human patient without cancer and/or non cancerous samples from the patient; in the methods for diagnosing or monitoring for metastasis, normal human control may also include samples from a human patient that is determined by reliable methods to have colon cancer which has not metastasized.

#### **Staging**

The invention also provides a method of staging colon cancer in a human patient.

The method comprises identifying a human patient having such cancer; analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG. Then, the method compares CSG levels in such cells, tissues, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of CSG is associated with a cancer which is regressing or in remission.

#### **Monitoring**

Further provided is a method of monitoring colon cancer in a human having such cancer for the onset of metastasis. The method comprises identifying a human patient having such cancer that is not known to have metastasized; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissue, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which has metastasized.

Further provided by this inventions is a method of monitoring the change in stage of colon cancer in a human having such cancer. The method comprises identifying a human patient having such cancer; periodically analyzing a sample

WO 99/60161

PCT/US99/10498

- 9 -

of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissue, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which is progressing in stage and a decrease in the levels of CSG is associated with a cancer which is regressing in stage or in remission.

Monitoring such patient for onset of metastasis is periodic and preferably done on a quarterly basis. However, this may be more or less frequent depending on the cancer, the particular patient, and the stage of the cancer.

#### **Assay Techniques**

Assay techniques that can be used to determine levels of gene expression, such as CSG of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, reverse transcriptase PCR (RT-PCR) assays, immunohistochemistry assays, in situ hybridization assays, competitive-binding assays, Western Blot analyses and ELISA assays. Among these, ELISAs are frequently preferred to diagnose a gene's expressed protein in biological fluids. An ELISA assay initially comprises preparing an antibody, if not readily available from a commercial source, specific to CSG, preferably a monoclonal antibody. In addition a reporter antibody generally is prepared which binds specifically to CSG. The reporter antibody is attached to a detectable reagent such as radioactive, fluorescent or enzymatic reagent, for example horseradish peroxidase enzyme or alkaline phosphatase.

To carry out the ELISA, antibody specific to CSG is incubated on a solid support, e.g., a polystyrene dish, that binds the antibody. Any free protein binding sites on the dish are then covered by incubating with a non-specific

WO 99/60161

PCT/US99/10498

- 10 -

protein such as bovine serum albumin. Next, the sample to be analyzed is incubated in the dish, during which time CSG binds to the specific antibody attached to the polystyrene dish. Unbound sample is washed out with buffer. A reporter antibody specifically directed to CSG and linked to horseradish peroxidase is placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to CSG. Unattached reporter antibody is then washed out. Reagents for peroxidase activity, including a colorimetric substrate are then added to the dish. Immobilized peroxidase, linked to CSG antibodies, produces a colored reaction product. The amount of color developed in a given time period is proportional to the amount of CSG protein present in the sample. Quantitative results typically are obtained by reference to a standard curve.

A competition assay may be employed wherein antibodies specific to CSG attached to a solid support and labeled CSG and a sample derived from the host are passed over the solid support and the amount of label detected attached to the solid support can be correlated to a quantity of CSG in the sample. Nucleic acid methods may be used to detect CSG mRNA as a marker for colon cancer. Polymerase chain reaction (PCR) and other nucleic acid methods, such as ligase chain reaction (LCR) and nucleic acid sequence based amplification (NASABA), can be used to detect malignant cells for diagnosis and monitoring of various malignancies. For example, reverse-transcriptase PCR (RT-PCR) is a powerful technique which can be used to detect the presence of a specific mRNA population in a complex mixture of thousands of other mRNA species. In RT-PCR, an mRNA species is first reverse transcribed to complementary DNA (cDNA) with use of the enzyme reverse transcriptase; the cDNA is then amplified as in a standard PCR reaction. RT-PCR can thus reveal by amplification the presence of a single species of mRNA. Accordingly, if the mRNA is highly specific for the cell that produces it, RT-PCR



WO 99/60161

PCT/US99/10498

- 11 -

can be used to identify the presence of a specific type of cell.

Hybridization to clones or oligonucleotides arrayed on a solid support (i.e., gridding) can be used to both detect the expression of and quantitate the level of expression of that gene. In this approach, a cDNA encoding the CSG gene is fixed to a substrate. The substrate may be of any suitable type including but not limited to glass, nitrocellulose, nylon or plastic. At least a portion of the DNA encoding the CSG gene is attached to the substrate and then incubated with the analyte, which may be RNA or a complementary DNA (cDNA) copy of the RNA, isolated from the tissue of interest.

Hybridization between the substrate bound DNA and the analyte can be detected and quantitated by several means including but not limited to radioactive labeling or fluorescence labeling of the analyte or a secondary molecule designed to detect the hybrid. Quantitation of the level of gene expression can be done by comparison of the intensity of the signal from the analyte compared with that determined from known standards. The standards can be obtained by *in vitro* transcription of the target gene, quantitating the yield, and then using that material to generate a standard curve.

The above tests can be carried out on samples derived from a variety of patients' cells, bodily fluids and/or tissue extracts (homogenates or solubilized tissue) such as from tissue biopsy and autopsy material. Bodily fluids useful in the present invention include blood, urine, saliva, or any other bodily secretion or derivative thereof. Blood can include whole blood, plasma, serum, or any derivative of blood.

#### EXAMPLES

The present invention is further described by the following examples. These examples are provided solely to illustrate the invention by reference to specific embodiments.

WO 99/60161

PCT/US99/10498

- 12 -

These exemplifications, while illustrating certain specific aspects of the invention, do not portray the limitations or circumscribe the scope of the disclosed invention.

WO 99/60161

PCT/US99/10498

- 13 -

**Example 1: CSGs**

Searches were carried out and CSGs identified using the following Search Tools as part of the LIFESEQ® database available from Incyte Pharmaceuticals, Palo Alto, CA:

1. Library Comparison (compares one library to one other library) allows the identification of clones expressed in tumor and absent or expressed at a lower level in normal tissue.

2. Subsetting is similar to library comparison but allows the identification of clones expressed in a pool of libraries and absent or expressed at a lower level in a second pool of libraries.

3. Transcript Imaging lists all of the clones in a single library or a pool of libraries based on abundance. Individual clones can then be examined using Electronic Northernblots to determine the tissue sources of their component ESTs.

4. Protein Function: Incyte has identified subsets of ESTs with a potential protein function based on homologies to known proteins. Some examples in this database include Transcription Factors and Proteases. We identified some leads by searching in this database for clones whose component ESTs showed disease specificity.

Electronic subtractions, transcript imaging and protein function searches were used to identify clones, whose component ESTs were exclusively or more frequently found in libraries from specific tumors. Individual candidate clones were examined in detail by checking where each EST originated.

WO 99/60161

PCT/US99/10498

- 14 -

Table 1: CSGs

SEQ ID	Clone ID #	Gene ID #	
NO:			
1	238330	242807	Transcript Imaging
2	1295234	239588	Subsetting
3	1341701	29634	Transcript Imaging
4	816257	233421	Subsetting
5	775133	245080	Subsetting
6	1335450	245811	Subsetting
7	2348122	233711	Transcript Imaging
8	3228674	230273	Subsetting
9	1632174	229022	Transcript Imaging

The following example was carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. Routine molecular biology techniques of the following example can be carried out as described in standard laboratory manuals, such as Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

**Example 2: Relative Quantitation of CSG Gene Expression**

Real-Time quantitative PCR with fluorescent Taqman probes is a quantitation detection system utilizing the 5'-3' nuclease activity of Taq DNA polymerase. The method uses an internal fluorescent oligonucleotide probe (Taqman) labeled with a 5' reporter dye and a downstream, 3' quencher dye. During PCR, the 5'-3' nuclease activity of Taq DNA polymerase releases the reporter, whose fluorescence can then be detected by the laser detector of the Model 7700 Sequence Detection System (PE Applied Biosystems, Foster City, CA, USA).

WO 99/60161

PCT/US99/10498

- 15 -

Amplification of an endogenous control is used to standardize the amount of sample RNA added to the reaction and normalize for Reverse Transcriptase (RT) efficiency. Either cyclophilin, glyceraldehyde-3-phosphate dehydrogenase (GAPDH) or 18S ribosomal RNA (rRNA) is used as this endogenous control. To calculate relative quantitation between all the samples studied, the target RNA levels for one sample are used as the basis for comparative results (calibrator). Quantitation relative to the "calibrator" can be obtained using the standard curve method or the comparative method (User Bulletin #2: ABI PRISM 7700 Sequence Detection System).

To evaluate the tissue distribution, and the level of CSGs in normal and tumor tissue, total RNA was extracted from normal tissues, tumor tissues, and from tumors and the corresponding matched normal tissues. Subsequently, first strand cDNA was prepared with reverse transcriptase and the polymerase chain reaction was done using primers and Taqman probe specific to the CSG. The results were analyzed using the ABI PRISM 7700 Sequence Detector. The absolute numbers are relative levels of expression of the CSG compared to the calibrator.

#### Comparative Examples

Similar mRNA expression analysis for genes coding for the diagnostic markers PSA (Prostate Specific Antigen) and PLA2 (Phospholipase A2) was performed for comparison. PSA is currently the only cancer screening marker available in clinical laboratories. When the panel of normal pooled tissues was analyzed, PSA was expressed at very high levels in prostate, with a very low expression in breast and testis. After analysis of more than 55 matching samples from 14 different tissues, the data corroborated the tissue specificity seen with normal tissue samples. PSA expression was compared in cancer and normal adjacent tissue for 12 matching samples of prostate tissue. The relative levels of PSA were higher in 10 cancer samples (83%). Clinical data

WO 99/60161

PCT/US99/10498

- 16 -

recently obtained support the utilization of PLA2 as a staging marker for late stages of prostate cancer. mRNA expression data described herein showed overexpression of the mRNA in 8 out of the 12 prostate matching samples analyzed (66%). PLA2 had high levels of mRNA expression in small intestine, prostate, liver, and pancreas.

**Measurement of SEQ ID NO:3; Clone ID 1341701; Gene ID 29634 (Cln106)**

Absolute numbers are depicted in Table 2 as relative levels of expression of Cln106 (SEQ ID NO:3) in 12 normal different tissues. All the values are compared to normal testis (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

**Table 2: Relative levels of Cln106 Expression in Pooled Samples**

Tissue	NORMAL
Colon-Ascending	110
Endometrium	0
Kidney	0
Liver	0
Ovary	0
Pancreas	0
Prostate	16
Small Intestine	0
Spleen	0
Stomach	0
Testis	1
Uterus	0

The relative levels of expression in Table 2 show for the CSG Cln106 (SEQ ID NO:3), mRNA expression is more than 6 fold higher in the pool of normal ascending colon (110) compared with prostate (16). Testis, the calibrator, with a relative expression level of 1, is the only other tissue expressing the mRNA for Cln106 (SEQ ID NO:3). These results demonstrate that mRNA expression of this CSG is highly specific for colon.

WO 99/60161

PCT/US99/10498

- 17 -

The absolute numbers in Table 2 were obtained analyzing pools of samples of a particular tissue from different individuals. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 3.

The absolute numbers in Table 3 are relative levels of expression of Cln106 (SEQ ID NO:3) in 57 pairs of matching samples. All the values are compared to normal testis (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 3: Relative levels of Cln106 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matching Normal Adjacent
Sto AC93	Stomach 1	4	96
Sto AC99	Stomach 2	0.4	0.5
Sml 21XA	Small Intestine 1	0	0
Sml H89	Small Intestine 2	0.93	1.28
Cln B56	Colon-Cecum(A) 1	317	101
Cln AS45	Colon-Ascending(A) 2	316.3	146.5
Cln CM67	Colon-Cecum(B) 3	481.0	217.5
Cln AS67	Colon-Ascending(B) 4	858.1	220.6
Cln AS43	Colon-Ascending(C) 5	1370	98
Cln AS46	Colon-Ascending(C) 6	3051	375
Cln AS98	Colon-Ascending(C) 7	26	42
Cln AS89	Colon-Ascending(D) 8	524.6	11.0
Cln TX01	Colon-Transverse(B) 9	2886.3	1992.0
Cln TX89	Colon-Transverse(B) 10	146.0	35.9
Cln TX67	Colon-Transverse(C) 11	2.9	421.7
Cln MT38	Colon-Splenic Flexure(M) 12	1681	187

WO 99/60161

PCT/US99/10498

- 18 -

Cln SG89	Colon-Sigmoid(B)13	1063.8	31.1
Cln SG67	Colon-Sigmoid(C)14	8.5	9.4
Cln SG33	Colon-Sigmoid(C)15	264	549
Cln SG45	Colon-Sigmoid(D)16	580.0	114.6
Cln B34	Colon-Rectosigmoid(A)17	97	244
Cln CXGA	Colon-Rectum(A)18	45.1	273.4
Cln RC67	Colon-Rectum(B)19	2.7	20.0
Cln C9XR	Colon-Rectosigmoid(C)20	609	460
Cln RS45	Colon-Rectosigmoid(C)21	472.8	144.0
Cln RC01	Colon-Rectum(C)22	568	129
Cln RC89	Colon-Rectum(D)23	4.6	322.91
Bld 46XK	Bladder 1	0.2	0
Bld 66X	Bladder 2	1	1
Bld 32XK	Bladder 3	0.0	0.0
Kid 126XD	Kidney 1	0	0
Kid 12XD	Kidney 2	0	0
Kid 5XD	Kidney 3	0.0	1.0
Kid 6XD	Kidney 4	0.0	0.0
Kid 106XD	Kidney 5	0.4	0.0
Liv 42X	Liver 1	0.0	0.0
Liv 15XA	Liver 2	0.0	0.0
Liv 94XA	Liver 3	0.0	0.0
Lng AC69	Lung 1	2	0
Lng BR94	Lung 2	0	0
Lng 47XQ	Lung 3	0	0
Mam 59X	Mammary Gland 1	0	0
Mam B011X	Mammary Gland 2	0	0
Mam A06X	Mammary Gland 3	0	0
Ovr 103X	Ovary 1	0.04	2.08
Ovr 130X	Ovary 2	0.1	2.76



WO 99/60161

PCT/US99/10498

- 19 -

Pan 71XL	Pancreas 1	4.08	0.1
Pan 82XP	Pancreas 2	0	0
Pro 12B	Prostate 1	0.3	0
Pro 23B	Prostate 2	3	4
Pro 13XB	Prostate 3	2	7
Pro 34B	Prostate 4	0.54	4.01
Pro 20XB	Prostate 5	4.8	4.3
Pro 65XB	Prostate 6	0.7	1.3
Tst 39X	Testis 1	2.78	0
End 8XA	Endometrium 1	0	0.2
Utr 85XU	Uterus 1	1.26	0

0= Negative

When matching samples were analyzed, the higher levels of expression were in the colon, showing a high degree of tissue specificity for this tissue. These results confirm the tissue specificity results obtained with the panel of normal pooled samples (Table 2). Furthermore, the level of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer stage (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 3 shows overexpression of Cln106 (SEQ ID NO:3) in 15 colon cancer tissues compared with their respective normal adjacent (colon samples #1, 2, 3, 4, 5, 6, 8, 9, 10, 12, 13, 16, 20, 21, and 22). There is overexpression in the cancer tissue for 65% of the colon matching samples tested (total of 23 colon matching samples). The matching sample Pan 71XL is a secondary cancer in pancreas, the primary cancer in that individual was a duodenal cancer.

Altogether, the high level of tissue specificity, plus the mRNA overexpression in 65% of the colon matching samples

WO 99/60161

PCT/US99/10498

- 20 -

tested are demonstrative of CSG Cln106 (SEQ ID NO:3) being a diagnostic marker for colon cancer.

Measurement of SEQ ID NO:4; Clone ID 816257; Gene ID 406452 (Cln107)

Absolute numbers as depicted in Table 4 are relative levels of expression of CSG Cln107 (SEQ ID NO:4) in 12 normal different tissues. All the values are compared to normal small intestine (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 4: Relative levels of Cln107 Expression in Pooled Samples

Tissue	NORMAL
Colon-Ascending	3.2
Endometrium	0
Kidney	0.2
Liver	0
Ovary	0
Pancreas	0
Prostate	0.1
Small Intestine	1
Spleen	0
Stomach	0.3
Testis	0
Uterus	0

The relative levels of expression in Table 4 show that mRNA expression of the CSG Cln107 (SEQ ID NO:4) is more than 10 fold higher in the pool of normal ascending colon (3.2), five fold higher in small intestine (1), and 1.5 fold higher in stomach (0.3), compared with the next higher expressor (0.2 for kidney). Seven of the pooled tissues samples analyzed were negative and prostate showed a relative expression of 0.1 for Cln107 (SEQ ID NO:4). These results demonstrate that Cln107 mRNA expression is highly specific for colon, small intestine, and in a lower degree for stomach.

The absolute numbers in Table 4 were obtained analyzing pools of samples of a particular tissue from different

WO 99/60161

PCT/US99/10498

- 21 -

individuals. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 5.

The absolute numbers in Table 5 are relative levels of expression of Cln107 (SEQ ID NO:4) in 57 pairs of matching samples. All the values are compared to normal small intestine (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

**Table 5: Relative levels of Cln107 Expression in Individual Samples**

Sample ID	Tissue	Cancer	Matching Normal Adjacent
Sto AC93	Stomach 1	8.9	13.4
Sto AC99	Stomach 2	6.0	0.9
Sml 21XA	Small Intestine 1	1.07	1.42
Sml H89	Small Intestine 2	0.97	4.13
Cln B56	Colon-Cecum(A) 1	2	16
Cln AS45	Colon-Ascending(A) 2	0.7	2.1
Cln CM67	Colon-Cecum(B) 3	1.6	2.1
Cln AS67	Colon-Ascending(B) 4	1.2	6.2
Cln AS43	Colon-Ascending(C) 5	13.5	0.5
Cln AS46	Colon-Ascending(C) 6	9.7	23.6
Cln AS98	Colon-Ascending(C) 7	28.1	1.4
Cln AS89	Colon-Ascending(D) 8	0.9	3.1
Cln TX01	Colon-Transverse(B) 9	3.0	10.6
Cln TX89	Colon-Transverse(B) 10	4.5	0.6
Cln TX67	Colon-Transverse(C) 11	3.6	3.4
Cln MT38	Colon-Splenic Flexure (M) 12	4.0	2.6
Cln SG89	Colon-Sigmoid(B) 13	4.7	0.9

WO 99/60161

PCT/US99/10498

- 22 -

Cln SG67	Colon-Sigmoid(C)14	1.0	1.3
Cln SG33	Colon-Sigmoid(C)15	14.2	7.6
Cln SG45	Colon-Sigmoid(D)16	4.8	6.0
Cln B34	Colon-Rectosigmoid(A)17	3	2
Cln CXGA	Colon-Rectum(A)18	4.4	1.9
Cln RC67	Colon-Rectum(B)19	0.1	0.4
Cln C9XR	Colon-Rectosigmoid(C)20	5	3
Cln RS45	Colon-Rectosigmoid(C)21	11.4	4.6
Cln RC01	Colon-Rectum(C)22	1.8	2.3
Cln RC89	Colon-Rectum(D)23	0.1	5.35
Bld 46XK	Bladder 1	0.2	0
Bld 66X	Bladder 2	1	1
Bld 32XK	Bladder 3	0.1	0.1
Kid 126XD	Kidney 1	0	0.02
Kid 12XD	Kidney 2	0.1	0.2
Kid 5XD	Kidney 3	0.3	0.0
Kid 6XD	Kidney 4	0.1	0.1
Kid 106XD	Kidney 5	0.0	0.1
Liv 42X	Liver 1	7.9	0.002
Liv 15XA	Liver 2	0.0	0.0
Liv 94XA	Liver 3	0.0	0.0
Lng AC69	Lung 1	1.6	0.2
Lng BR94	Lung 2	0.4	0
Lng 47XQ	Lung 3	0.78	0.2
Mam 59X	Mammary Gland 1	0.05	0.3
Mam B011X	Mammary Gland 2	0.01	0.004
Mam A06X	Mammary Gland 3	0.22	0
Ovr 103X	Ovary 1	0.01	0.01
Ovr 130X	Ovary 2	0.09	0.1
Pan 71XL	Pancreas 1	2.51	2.81

WO 99/60161

PCT/US99/10498

- 23 -

Pan 82XP	Pancreas 2	0	0.62
Pro 12B	Prostate 1	0.3	0.1
Pro 23B	Prostate 2	0.3	0.2
Pro 13XB	Prostate 3	0	0
Pro 34B	Prostate 4	0.04	0.22
Pro 20XB	Prostate 5	0.4	0.1
Pro 65XB	Prostate 6	0.0	0.1
Tst 39X	Testis 1	0.02	0.01
End 8XA	Endometrium 1	0.01	0.5
Utr 85XU	Uterus 1	0.03	0

0= Negative

When matching samples were analyzed, the higher levels of expression were in colon, stomach, and small intestine, showing a high degree of tissue specificity for colon tissues. These results confirm the tissue specificity results obtained with normal pooled samples (Table 4). Furthermore, the level of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer stage (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 5 shows overexpression of Cln107 (SEQ ID NO:4) in 11 colon cancer tissues compared with their respective normal adjacent (colon samples #5, 7, 10, 11, 12, 13, 15, 17, 18, 20, and 21). There is overexpression in the cancer tissue for 48% of the colon matching samples tested (total of 23 colon matching samples). The matching sample Pan 71XL is a secondary cancer in pancreas, the primary cancer in that individual was a duodenal cancer.

Altogether, the high level of tissue specificity, plus the mRNA overexpression in almost half of the colon, stomach, and small intestine matching samples tested are demonstrative

WO-99/60161

PCT/US99/10498

- 24 -

of CSG Cln107 (SEQ ID NO:4) being a diagnostic marker for colon cancer.

Measurement of SEQ ID NO:5; Clone ID 775133; Gene ID 24508 (Cln108)

The absolute numbers shown in Table 6 are relative levels of expression of CSG Cln108 (SEQ ID NO:5) in 12 normal different tissues. All the values are compared to normal small intestine (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 6: Relative levels of Cln108 Expression in Pooled Samples

Tissue	NORMAL
Colon-Ascending	2846.5
Endometrium	1
Kidney	5.5
Liver	18.7
Ovary	3.4
Pancreas	198.1
Prostate	1024
Small Intestine	810.8
Spleen	32.2
Stomach	9981.2
Testis	0
Uterus	294.1

The relative levels of expression in Table 6 show that mRNA expression of CSG Cln108 (SEQ ID NO:5) is more than 10 fold higher in the pool of normal ascending colon (2846.5) and almost ten fold higher in stomach (9981.2), compared to the expression level in any other tissue analyzed. These results demonstrate that mRNA expression of this CSG is also highly specific for colon and stomach.

The absolute numbers in Table 6 were obtained analyzing pools of samples of a particular tissue from different individuals. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 7.

WO 99/60161

PCT/US99/10498

- 25 -

The absolute numbers depicted in Table 7 are relative levels of expression of Cln108 (SEQ ID NO:5) in 57 pairs of matching samples. All the values are compared to normal small intestine (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 7: Relative levels of Cln108 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matching Normal Adjacent
Sto AC93	Stomach 1	28696	34842
Sto AC99	Stomach 2	21523	30862
Sml 21XA	Small Intestine 1	2944	964.4
Sml H89	Small Intestine 2	244.5	3513.2
Cln B56	Colon-Cecum(A) 1	27242	24637
Cln AS45	Colon-Ascending(A) 2	5827.0	8771.0
Cln CM67	Colon-Cecum(B) 3	4251.0	4684.0
Cln AS67	Colon-Ascending(B) 4	564.0	1949.0
Cln AS43	Colon-Ascending(C) 5	50310	10949
Cln AS46	Colon-Ascending(C) 6	246044	120073
Cln AS98	Colon-Ascending(C) 7	40442	17482
Cln AS89	Colon-Ascending(D) 8	5730.0	1581.0
Cln TX01	Colon-Transverse(B) 9	22281.0	114784.0
Cln TX89	Colon-Transverse(B) 10	11026.0	1639.0
Cln TX67	Colon-Transverse(C) 11	17004.0	11654.0
Cln MT38	Colon-Splenic Flexure(M) 12	77589	31620
Cln SG89	Colon-Sigmoid(B) 13	140339.0	49617.0
Cln SG67	Colon-Sigmoid(C) 14	4951.0	7905.0
Cln SG33	Colon-Sigmoid(C) 15	60875	120490

WO 99/60161

PCT/US99/10498

- 26 -

Cln SG45	Colon-Sigmoid(D) 16	30437.0	47267.0
Cln B34	Colon-Rectosigmoid(A) 17	5848	5861
Cln CXGA	Colon-Rectum(A) 18	13877.0	9787.0
Cln RC67	Colon-Rectum(B) 19	1703.0	26589.0
Cln C9XR	Colon-Rectosigmoid(C) 20	2458	19071
Cln RS45	Colon-Rectosigmoid(C) 21	95523	61939
Cln RC01	Colon-Rectum(C) 22	98891.0	80047.0
Cln RC89	Colon-Rectum(D) 23	17.0	1775
Bld 46XK	Bladder 1	0	8
Bld 66X	Bladder 2	397	44
Bld 32XK	Bladder 3	0.0	16.0
Kid 126XD	Kidney 1	32	22
Kid 12XD	Kidney 2	6	0
Kid 106XD	Kidney 3	4.0	33.0
Liv 42X	Liver 1	4783	0
Liv 15XA	Liver 2	4.0	10.0
Liv 94XA	Liver 3	159.0	21.0
Lng AC69	Lung 1	222	295
Lng BR94	Lung 2	112	0
Lng 47XQ	Lung 3	30	69
Lng AC66	Lung 4	29	137
Mam 59X	Mammary Gland 1	56	0
Mam B011X	Mammary Gland 2	54	31
Mam A06X	Mammary Gland 3	12	0
Ovr 103X	Ovary 1	37	0
Pan 71XL	Pancreas 1	13203	4163
Pan 82XP	Pancreas 2	39.1	0
Pro 12B	Prostate 1	386	88
Pro 23B	Prostate 2	250	23
Pro 13XB	Prostate 3	92	731



WO 99/60161

PCT/US99/10498

- 27 -

Pro 34B	Prostate 4	33.3	265.7
Pro 20XB	Prostate 5	454.6	1908.9
Pro 65XB	Prostate 6	733.5	922.0
End 8XA	Endometrium 1	5	92
Utr 85XU	Uterus 1	98.9	21.8
Utr 23XU	Uterus 2	35.3	0
Utr 135X0	Uterus 3	39.2	43.8
Utr 141X0	Uterus 4	212.1	55.9

0= Negative

When matching samples were analyzed, the higher levels of expression were in colon and stomach, showing a high degree of tissue specificity for these two tissues. These results confirm the tissue specificity results obtained with normal pooled samples (Table 6). Furthermore, the level of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer stage (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 7 shows overexpression of CSG Cln108 (SEQ ID NO:5) in 13 colon cancer tissues compared with their respective normal adjacent (colon samples #1, 5, 6, 7, 8, 9, 10, 11, 12, 13, 18, 21, and 22). There is overexpression in the cancer tissue for 56% of the colon matching samples tested (total of 23 colon matching samples). The matching sample Pan 71XL is a secondary cancer in pancreas, the primary cancer in that individual was a duodenal cancer.

Altogether, the high level of tissue specificity, plus the mRNA overexpression in more than half of the colon, stomach, and small intestine matching samples tested are demonstrative of this CSG, Cln108 (SEQ ID NO:5), also being a diagnostic marker for colon cancer.

WO 99/60161

PCT/US99/10498

- 28 -

Measurement of SEQ ID NO:7; Clone ID 2348122; Gene ID 23371  
(Cln109)

The absolute numbers depicted in Table 8 are relative levels of expression of CSG Cln109 (SEQ ID NO:7) in 12 normal different tissues. All the values are compared to normal ovary (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 8: Relative levels of Cln109 Expression in Pooled Samples

Tissue	NORMAL
Colon-Ascending	28.8
Endometrium	0.45
Kidney	0.41
Liver	0.72
Ovary	0.07
Pancreas	82.8
Prostate	124.3
Small Intestine	626.4
Spleen	1.2
Stomach	12.05
Testis	1.51
Uterus	52.99

The relative levels of expression in Table 8 show that mRNA expression of CSG Cln109 (SEQ ID NO:7), is more than 5 fold higher in the pool of normal small intestine (626.4) compared to the expression level in any other tissue analyzed. These results demonstrate that Cln109 (SEQ ID NO:7) mRNA expression is highly specific for small intestine

The absolute numbers in Table 8 were obtained analyzing pools of samples of a particular tissue from different individuals. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 9.

The absolute numbers depicted in Table 9 are relative levels of expression of Cln109 (SEQ ID NO:7) in 53 pairs of matching samples. All the values are compared to normal ovary (calibrator). A matching pair is formed by mRNA from the

WO 99/60161

PCT/US99/10498

- 29 -

cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 9: Relative levels of Cln109 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matching Normal Adjacent
Sto AC93	Stomach 1	2574	1310
Sto AC99	Stomach 2	4153	5
Sml 21XA	Small Intestine 1	2667	13663.8
Sml H89	Small Intestine 2	57.8	904.29
Cln B56	Colon-Cecum(A) 1	6794	299
Cln AS45	Colon-Ascending(A) 2	814.6	105.8
Cln CM67	Colon-Cecum(B) 3	294.6	36.1
Cln AS67	Colon-Ascending(B) 4	2.2	26.3
Cln AS43	Colon-Ascending(C) 5	111	377
Cln AS46	Colon-Ascending(C) 6	1180	352
Cln AS98	Colon-Ascending(C) 7	1075	92
Cln AS89	Colon-Ascending(D) 8	14022.7	87.5
Cln TX01	Colon-Transverse(B) 9	1027.6	282.1
Cln TX89	Colon-Transverse(B) 10	2.5	23.7
Cln TX67	Colon-Transverse(C) 11	0.1	72.3
Cln MT38	Colon-Splenic Flexure(M) 12	372	88
Cln SG89	Colon-Sigmoid(B) 13	179.2	33.4
Cln SG67	Colon-Sigmoid(C) 14	85.0	94.7
Cln SG33	Colon-Sigmoid(C) 15	5461	377
Cln SG45	Colon-Sigmoid(D) 16	762.7	15.9
Cln B34	Colon-Rectosigmoid(A) 17	460	1
Cln RC67	Colon-Rectum(B) 18	64.5	136.2
Cln C9XR	Colon-Rectosigmoid(C) 19	441	34
Cln RS45	Colon-Rectosigmoid(C) 20	1931	195

WO 99/60161

PCT/US99/10498

- 30 -

Cln RC01	Colon-Rectum(C) 21	72.8	19.1
Cln RC89	Colon-Rectum(D) 22	4.8	90.2
Bld 46XK	Bladder 1	4	3
Bld 66X	Bladder 2	1	0
Bld 32XK	Bladder 3	0.1	307.6
Kid 126XD	Kidney 1	0	2
Kid 12XD	Kidney 2	3	16
Kid 5XD	Kidney 3	0.0	0.3
Kid 6XD	Kidney 4	18.5	1.2
Liv 42X	Liver 1	21	0.03
Liv 15XA	Liver 2	0.5	0.4
Liv 94XA	Liver 3	0.4	0.0
Lng AC69	Lung 1	0.1	0
Lng BR94	Lung 2	3	0
Lng 60XL	Lung 3	0.1	0
Mam 59X	Mammary Gland 1	0	4
Mam B011X	Mammary Gland 2	8	13
Mam A06X	Mammary Gland 3	4.7	9.6
Pan 71XL	Pancreas 1	8902.5	1428.2
Pan 82XP	Pancreas 2	0.2	9.3
Pro 12B	Prostate 1	9	20
Pro 23B	Prostate 2	191	88
Pro 13XB	Prostate 3	12	460
Pro 34B	Prostate 4	3.2	80.4
Tst 39X	Testis 1	29.9	0
End 8XA	Endometrium 1	0.3	21
Utr 85XU	Uterus 1	244.7	592.2
Ovr 63A	Ovary 1	11.4	0
Ovr A1C	Ovary 2	68.4	0

0= Negative

WO 99/60161

PCT/US99/10498

- 31 -

When matching samples were analyzed, the higher levels of expression were in small intestine, colon and stomach, showing a high degree of tissue specificity for these three colon tissues. These results confirm the tissue specificity results obtained with normal pooled samples for small intestine (Table 8). Furthermore, the level of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer stage (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 9 shows overexpression of CSG, Cln109 (SEQ ID NO:7) in 15 colon cancer tissues compared with their respective normal adjacent (colon samples #1, 2, 3, 6, 7, 8, 9, 12, 13, 15, 16, 17, 19, 20, and 21). There is overexpression in the cancer tissue for 68% of the colon matching samples tested (total of 22 colon matching samples). The matching sample Pan 71XL is a secondary cancer in pancreas, the primary cancer in that individual was a duodenal cancer.

Altogether, the high level of tissue specificity, plus the mRNA overexpression in more than half of the colon, stomach, and small intestine matching samples tested are demonstrative of CSG Cln109 (SEQ ID NO:7) being a diagnostic marker for colon cancer. The amino acid sequence encoded by the open reading frame of Cln109 is depicted in SEQ ID NO:10.

WO 99/60161

PCT/US99/10498

- 32 -

What is claimed is:

1. A method for diagnosing the presence of colon cancer in a patient comprising:

(a) measuring levels of CSG in a sample of cells, tissue or bodily fluid obtained from the patient; and

(b) comparing the measured levels of CSG with levels of CSG in a sample of cells, tissue or bodily fluid obtained from a control, wherein an increase in measured levels of CSG in the patient versus the CSG levels in the control is associated with the presence of colon cancer.

2. A method of diagnosing metastatic colon cancer in a patient comprising:

(a) measuring levels of CSG in a sample of cells, tissue, or bodily fluid obtained from the patient; and

(b) comparing the measured levels of CSG with levels of CSG in a sample of cells, tissue, or bodily fluid obtained from a control, wherein an increase in measured CSG levels in the patient versus the CSG levels in the control is associated with a cancer which has metastasized.

3. A method of staging colon cancer in a patient comprising:

(a) identifying a patient suffering from colon cancer;

(b) measuring levels of CSG in a sample of cells, tissue, or bodily fluid obtained from the patient; and

(c) comparing the measured levels of CSG with levels of CSG in a sample of cells, tissue, or bodily fluid obtained from a control, wherein an increase in the measured levels of CSG versus the levels of CSG in the control is associated with a cancer which is progressing and a decrease in the measured levels of CSG versus the levels of CSG in the control is associated with a cancer which is regressing or in remission.

WO 99/60161

PCT/US99/10498

- 33 -

4. A method of monitoring colon cancer in a patient for the onset of metastasis comprising:

(a) identifying a patient having colon cancer that is not known to have metastasized;

(b) periodically measuring CSG levels in samples of cells, tissue, or bodily fluid obtained from the patient; and

(c) comparing the periodically measured levels of CSG with levels of CSG in cells, tissue, or bodily fluid obtained from a control, wherein an increase in any one of the periodically measured levels of CSG in the patient versus the levels of CSG in the control is associated with a cancer which has metastasized.

5. A method of monitoring changes in a stage of colon cancer in a patient comprising:

(a) identifying a patient having colon cancer;

(b) periodically measuring levels of CSG in samples of cells, tissue, or bodily fluid obtained from the patient; and

(c) comparing the measured levels of CSG with levels of CSG in a sample of the same cells, tissue, or bodily fluid of a control, wherein an increase in any one of the periodically measured levels of CSG versus levels of CSG in the control is associated with a cancer which is progressing in stage and a decrease in any one of the periodically measured levels of CSG versus the levels of CSG in the control is associated with a cancer which is regressing in stage or in remission.

6. The method of claim 1, 2, 3, 4 or 5 wherein the CSG comprises SEQ ID NO:3, 4, 5 or 7.

WO 99/60161

PCT/US99/10498

## SEQUENCE LISTING

<110> Macina, Roberto A.  
Yang, Fei  
Sun, Yongming

<120> A Novel Method of Diagnosing, Monitoring and Staging  
Colon Cancers

<130> DEX-0035

<140>

<141>

<150> 60/086.266

<151> 1998-05-21

<160> 10

<170> PatentIn Ver. 2.0

<210> 1

<211> 487

<212> DNA

<213> Homo sapiens

<400> 1

```
tctgcctctg gccctcccag tgcacctgtt caatcccagc yccctccctga cctgtacaaa 60
tacacctgag gaccggctcg agcccagact tcttgccctt gctctgcact ctccaggtatt 120
ccctgctctt actccaaaaa gatggaccga ggtccgaagg ggcactgcca ctgtgggggg 180
catggccatc ctccaggtca ctgcggggtga acccctggcc atggcccagg gccctgcggg 240
ccaccccttg gccatggccc agggccctgc gggcaacccc ctggccatgg ccagggccc 300
tgggggcttc cccctggcca tggcccaggt caccacccc ctggtccaca tccctgagga 360
agtagaagaa aacaggacac aagatggcaa gcttgagaga attgcccagc tgacctggaa 420
tgaggcctaa accacaatct tctcttccta ataaacagcc tccctagaggc cacattctat 480
tcttttaa
```

<210> 2

<211> 739

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (693)

<220>

<221> unsure



WO 99/60161

PCT/US99/10498

&lt;222&gt; (695) .. (699)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (703) .. (705)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (708)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (710) .. (716)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (718)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (723) .. (726)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (728)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (732)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (737)

&lt;400&gt; 2

```
tctgaaactg tcagttccac cagcactgct tggatacttg taagtttcca gggggctgct 60
ttgcactcga aactgtcagc cccagaatgt tgacagtcgc tctcttagcc ctctctcttg 120
cctcagcctc tggcaatgcc attcaggtcca ggtcttccct ctatagtgga gagtatggaa 180
gtgggtgctg aaagcgatcc tctcattctg gcaaccagtt ggacggcccc atcaccgccc 240
tccgggtccg agtcaacaaa tactacatcg taggtcttca ggtgcgctat ggcaaggtgc 300
ggagcgacta tgtgggtggt cgcaacggag acctggagga gactttcttg caccctgggg 360
aatcagtcat ccaggttttc ggggaagtaca agtggtagct gaaqaagctg gtatttctga 420
cagacaaggg ccgctatctg tcttttggga sagacagtgg cacaagtttc aatgccgtcc 480
ccttgcaccc caacaccgtg ctccgcttca tcagtggccg gtcgggttct ctcatcgarg 540
ccattggcct gcaactggga gtttacccca ctagctgcag cagatgctga gctcctctc 600
cttggcaggg gcaactgtgt gaggagtaag aactccctta tcaactaacc ccattccaaat 660
ggctcaataa aaaaatatgg ttaaggttaa aaaaaaang gannnaanan nnnnnntnce 720
aannnnanct cncctgnta 739
```

WO 99/60161

PCT/US99/10498

<210> 3  
<211> 428  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (391)

<400> 3  
aattgtccgg ggtcgaataag aggagagcat gaatgagagt catccctcgca agtgtgcaga 60  
gtcttttgaag atgtggcattg atcgtgacct ccactgtagg cgccttaagt ttgaaggcca 120  
tccccctgag tcttggaagt ggacccctgc accggtcatt ctttatatct gtgaaggcat 180  
cctccgggttt taccttcttc agcagaaggt tgtgattacc aacgttgtta tgcacccatc 240  
caaagttttg gaattgcaga tgaacaagcg tggcttcagc atggaagtgg ggcagtatac 300  
ctttgttaac tgcctcttca tctctctcct gggaaaggca tctcttact ttgacctctg 360  
ctccagagga agatttcttc ttcattcata tncgagcagc agggcgacctg acagaaaatc 420  
tataaggcg 428

<210> 4  
<211> 1347  
<212> DNA  
<213> Homo sapiens

<400> 4  
ggaaaacccc tgagcacaaa gcaagaggca tcgaagcccc ctccgggagc cccgcaagcc 60  
aacagggggtg tctgtgccttg gtagtacttc cgcctgcgtc ctctgctggtc caggggccca 120  
gacgagcccc agcagggccca agtcccccat gtctggggct gggaggtggc tggggccctt 180  
gcaactgaggc tgcagaaagt ccagtcacct gatctgctgg aaaggagagag ggagagtgctc 240  
ctgcgcgggg agcaagaggt ggagagggag cggagaaatg ctctcttccc agaggctcttc 300  
tccccaacgc cagatgagaa ctctgacctg aactccaggc ctctcttccc ggcactccggc 360  
atcacggggc gttactcctg gtctgagctc cctctcttca gccccatcca cctacactca 420  
aacgtggcgt ggaacagtga agatccagtg gacagtgtc ctcccgggca gagaaagaag 480  
gagcaatggc acgttggcat caacccctcg gacggtatca actcagaggc cctgggaagcc 540  
atacgggtga cccgtcacaa gaacggcatg gcagagctgt gggaaatccg catctacgct 600  
agtgaggagg atgactgagc ctcgggatgg ggcgcccacc cctgcccctg ccttgacctt 660  
cgtggggaact gccaaagacc tgcgcaagcc cccaccctag gaaatgggtc ctagggtccag 720  
gatccaagaa ccacagctca tctgccaaca atcccaccat gggcacattt gggactgttg 780  
ggtttttcgt ttcctgttct atcttctctt agaaatgttt ctgcttctgg ggtctaaaag 840  
ttttggggat gaaatgggga cccctgctga ttcttctctg tcttaagact tggccaatg 900  
ccctgggtct aagaaagaa gagaccgct cctccacttt cagggtgaat tggcttccgc 960  
tagtctgagg gcagggggat cgggtcaaaga ggggtggaca gatcgagca ccttgagggg 1020  
ctgcgggtct gagggagga acactcagct cctccctctg agaagtccca agctgagagg 1080  
ggagacctgc ccttctccaa ccttgggaaa ccatccagtc tggggaggga ggcacaaatc 1140  
ccagtgtgga ggggtccctg gcagccctca aaccttctac cttgggtgac ccagccacac 1200  
ctgggtggaca caaagctctt acatcgtatg gatcccatga ggtatggctcc cttcacctgg 1260  
ggaaaaagtg acccagttta ggaactggag ggggggtctt gtcctccacc cctaaaactg 1320

WO 99/60161

PCT/US99/10498

ccctgaaataa acctggagtg agctggc

1347

&lt;210&gt; 5

&lt;211&gt; 1249

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (1034)..(1046)

&lt;400&gt; 5

ggcagagcct ggcagagcct ggcagagcct ggcagagcct ggcagagcct 60  
acccctctggg cccactgcat ctgagagagc ggcctctctg aggcacactac cccctcagca 120  
actgggaggt gggactgcta gaagctgggc cagggtgctg gtcagctgag tcaggagcct 180  
aacggcaccct ggcctgggac accctgacct ctccatcgaa gcaggggagc tgggagcctc 240  
gagccctcgg ggcctgggac accctcagcc accctcagcc tggacagagc gagagtgcta 300  
ggctgctgctt ggcctcctctt cctcactctt ggcctcagca cgcactcagc gcttattcga 360  
agctacatga ggcctcagca gaaacacctc cgtctgcccac gctgctgctg cctgcccacc 420  
aaggagatcc agctcagca ggcctcagca ggcctcagca ggcctcagca agccaactac 480  
tttgcgtctt aactcctgag tgggctcggc aacgtcgtgg ggcctcagca gctgcttgaa 540  
gaccgcatga tcatgagctc tctgaaaaac aatgtgggca gaggcctaaa catcgccctg 600  
gtgaatgaa ccacgggagc tctgctggga cagaagtcac ttgacatgca ctctggagat 660  
gttatgcacc tagtgaaact ccttaaaaga attccggggg gtgcaactgt gctggtggcc 720  
tctacagacg atccagggac caaatgaac gatgaaagca ggaactctt ctctgacttg 780  
gggagttctt acgcaaaaca actgggcttc cgggacagct gggctctcat agggagcaaa 840  
gacctcaggg gtcagagcct ccttgagcag ttctcagaga acagcccaga cacaacacaa 900  
tacgagggat ggcctcagca gctggagatg gagggtgcta tgcccgcgaa gccattttag 960  
ggtggtctgt ggcctcagca agccaggggc ctgaaagagc tctgctctga cttaggagtc 1020  
agagcccggc agcnnnnnnn nnnnnnnnnn nnnnnnnnnn gctggaagc tctgctgagt 1080  
ccttgacagc tctgctcggc ctctcctcct cggaaacaga accctcccac agcacatctt 1140  
acccggaaga ccaactcagc agggctcttc tggaaaccag tctctgctga gagaatgggg 1200  
tgctttcagc agggactgct gacggctggt cctgaggaag gacaaactc 1240

&lt;210&gt; 6

&lt;211&gt; 1220

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 6

gctttctgca cctcattcca cctcagagc gctttctggg aaagctgcat tctgttgagc 60  
tccagggcgc agtggaggga gggagtgagc gagctctctg taccacagga aagtgcagct 120  
gagactcaga caagattaca atgaaccaac tcagcttctt gctgttctct atagcgacca 180  
ccagagagat gagtacagat gaggtcaca cttacttcaa ggaatggacc tgttcttctg 240  
ctccatctct gcccagagag tgcaggaaga tcagagagat atgtcctagt gcatttgatg 300  
gctctgattt tctcggcact gagaatgggt tctcttacca gacctctctg gacatgacct 360  
ctgggggtgg cggcgggacc ctggtggcca gctgacagc gaatgacatg cgtgggagat 420  
gcacggtggg cgtcggctct tccagtcagc agggcagcaa agcagctctc ccagaggggg 480

WO 99/60161

PCT/US99/10498

acggcaactg ggcacactac aacacccctg gatctgcaga ggcggccacg agcgatgact 540  
acaagaaccc tggctactac gacatccagg ccaaggacct ggcctctcgg cacgtgcccc 600  
ataagtcccc catgcagcac tggagaaaca gctccctgct gaggctaccg acggacactg 660  
gcttccctca gacactggga cataactcgt ttggcatcta ccagaaatac ccagtgaaa 720  
atggagaagg aaagtctgg actgacaacg qcccggtgat cccctgtggt tatgattttg 780  
gcgacgcgcc qaataacagc tcttattact caccctatgg caacgggga ttactgagg 840  
gattcttcca gtccagggtt tttaataacg agagagcagc caacgccttg tctgctgga 900  
tggaggctcac cggatgtaac actgagcacc actgcattgg tggagcagga tactttccag 960  
aggccagttc ccagcagctg ggcctttttt ctggttttga ttggagtcca ttgggaacct 1020  
atgtctggtt cagcagcagc cgtgagataa ctgagggcagc tgtgcttcta ttctatcgt 1080  
gagagttttg tgggagggga cccagacctc tctcccaac catgagatcc caaggatgga 1140  
gaacaactta cccagtagct agaatgttaa tggcagaaga gaaaacataa aatcatattg 1200  
actcaaaaaa aaaaaaaaag 1220

&lt;210&gt; 7

&lt;211&gt; 2796

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 7

cggctcaggg gacaggaatg ggcacggcct ctcatcttcc ctacgacctc tgttcttccc 60  
tggccaagct ccaggggatt tgggggatgt gggaccccca atccccagcc cgggcttcag 120  
ccctttcccc ggtgttgaat ccagctccag ctccagctcc agctccaggt cgggcttcag 180  
ctccagccgc agcttaggca gctgaggttc tgtgtccag ttgttttcca atttccaccg 240  
ctccgtggat gaccttggga cctgcccagc cctgtttccc ctgcccagca ccacctttcc 300  
cgtggacaga gtggaaacct tggaaattcac agctcatgtt ctctctcaga agtttgagaa 360  
agaactttcc aaagttaggg aatatgtcca attaatagc gtgtatgaaa agaaactgtt 420  
aaacctaaat gtccgaattg acatcatgga gaaggatacc atttcttaca ctgaacttga 480  
cttcgagctg atcaaggtag aagtgaaggga gatggaaaaa ctggtcatat agctgaagg 540  
gagttttggg ggaagctcag aaattgttga ccagctggag gtggagataa gaaatatgac 600  
tctcttggtt gagaagcttg agacactaga caaaaaaat gtcttggcca ttgcccagga 660  
aatctggtct ctgaagacaa agctgaaggg gtgtgagggc tctaaagatc aaaaaccccc 720  
tgtctgtccc ccttttcccc ctccaggagg ctgtggtcal ggtggtgtgg tgaacatcac 780  
caaacctgct gtggttccgc tcaactggag agggttttct tatctatatg gtgcttgggg 840  
tagggattac tctccccagc atccaaaaca aggaactgtat tgggtggcgc cattgaatac 900  
agatgggagc ctgttggagt attatagact gtacacacac ctggatgatc tgcatttga 960  
tataaatgct cagagattgc ggaacacctt tggccaaggc agtgggtacg cagtttaca 1020  
caacaacatg tacttcaaca tgtacaacac cgggaataat gccagagtta acctgaccac 1080  
caacacgatt gctgtgacct aaactctccc taatgtgtcc tataataacc gcttttcata 1140  
tgttaattgt gcttggcagg atattgactt tgcgtggat gagaatgga ttgtgggtat 1200  
ttattcaact gaagccagca ctggtaacat ggtgattagt aaactcaatg acaccacact 1260  
tcagggtgctt aacatttggc ataccaagca gtataaacca tctgcttcta acgacctcat 1320  
ggtatgtggg gtctgttatg ccacctgtac tatgaacacc agaacagaaq agatttttta 1380  
ctattatgac acaaacacag ggaasgaggg caaactagac attgtaatgc ataagatgca 1440  
ggaaaaagtg cagagcattt actataaccc ttttgaccag aaactttatg tctataacga 1500  
tggttacctt ctgaatttat atcttctctg ctgacagaaq cccagtaag ctgttttaga 1560  
gttaggttga aagagaaaaa gtttcttga aaaaatagtc tctccactta cttagatatc 1620  
tgcaggggtg tctaaaaagt tgttcatctt gcagcaatgt ctagggtgat agttctacca 1680

WO 99/60161

PCT/US99/10498

cactagagat ctaggacatt tgccttgatt tggtagatc tcttgggaat catctgcctc 1740  
ttcaggcgca ttttgcaata aagctctgtc agggtaggat tgcagaggt ctaggggcac 1800  
tgtgggccta gtgaagccta ctgtgaggag gcttcctag aagccttaaa ttagggaatta 1860  
aggaaactaa aactcagtat ggcgtctagg gattctttgt acaggaaata ttgccaatg 1920  
accagtcctc atccatgtag caccactaat tcttccatgc ctggaagaaa cctgggggac 1980  
tagttaggta gattaatata tggagctccg cgagggacca aatctccaac ttttttttc 2040  
cctcactaca cctggaatga tgccttgat gtggcagata agtaaatttg gcattgctat 2100  
atattctaca tctgtaaagt gctgagttt atggagagag gcctttttat gcattaaatt 2160  
gtacatggca aacaaatccc agaggatct gtagatgagg cacttgctt tcttttttc 2220  
tcattgtcca ccttactaaa agtcagtaga atcttctacc tcataacttc cttccaaagg 2280  
cagctcagaa gattagaacc agacttaata accaatcca ccccccacca accccttct 2340  
actgcctact ttaaaaaaat taatagttt ctatggaaat gatctaagat tagaaaaatt 2400  
aattttctt aatttcatta tggactttt ttacatgac tctaaagacta taagasaatt 2460  
tgatggcag gacaaaagt tgcactttat tgttatctaa taaagactt ggagcatatg 2520  
tgcaacttat gagtgtatca gtgttgcat graatttttg cctttgttca agcctgggaa 2580  
ttgttaagaa atgaataatt aattttttt tctaggacga gctatagaaa agctattgag 2640  
agtatctagt taactagtc agtaattgga aacctgtcg gtgtatgta tctgcttctg 2700  
tgcctttgaa tgaatttat atctagctt tgcctgttt tctttctatg ttaagttct 2760  
agtctatagg atggcagtt taaatgtctt actccc 2792

&lt;210&gt; B

&lt;211&gt; 2331

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (675)

&lt;400&gt; B

tttatcacgg gctcaactgc aacaaaaaac ttccttgaca gctccacaaa cttagggcac 60  
agttaggaat caacaaatat ccacagcagc ccagatgcaa gtggaacaa accctcatct 120  
gctcaactca caacctcagg tctgtgagaa tctacaacct cactcatcag tccaggctca 180  
actgaaataa caacgttacc tggcagtacc acaacaccag gcctcagtag ggcatctacc 240  
acctcttaca gtgcccag atcaccagac caaacactct cactgcccag catgagaago 300  
tccagcatca gtggagaacc caccagcttg tatagccaag cagagtcac acacacaaca 360  
gcgttccctg ccagcacac caccacaggc cttagtcagg aatcaacaa tttccatagt 420  
aagccagggt caactgagac aacactgtcc cctggcagca tcacaacttc atcttttgc 480  
caagaattta ccacctctca tagccaacca ggctcagctc tctcaacagt gtcacctgcc 540  
agcaccacag tgcacaggct tagtgaggaa tctaccacct tctacagcag cctaggctca 600  
actgaaacca cagcgttttc tcacagcaac acaatgtcca tctatagtca acaatctaca 660  
cccttccctg acagncagg cttcactcac acagtgttac ctgccacct cacaaccaca 720  
gacattggtc aggaatcaac agccttccac agcagctcag acgcaactgg aacaaacccc 780  
ttacctgccc gctccacagc cttagacctt gttggagaac ctacaacttt ctacatcagc 840  
ccatccctca ctacacaaac actctttcct gcgagtcca gcacatcagg cctcactgag 900  
gaatctacca ccttccacac cagtccaagc ttcacttcta caattgtgc tactgaaagc 960  
ctggaaaacct tagcaccagg gtctgtccag gaaggacaaa tttggaatgg aaaaacaatg 1020  
gtctgtcccc aaggctacgt tggttaccag tgcctgtccc ctctggaatc cttccctgta 1080

WO 99/60161

PCT/US99/10498

gaaaccccg g aaaaactcaa cggccacttta ggtatgacag tgaagtgac ttacagaaat 1140  
ttcacagaaa agatgaatga cgcattctcc caggaatacc agaacttcag taccctcttc 1200  
aagaatcggg tggatgctgt ttggaagggc gacaaatctc ctcaatatac aggggtgaa 1260  
attcggagat tgcctaacgg tagcatcgtg gtcaagaacg atgtcatcct ggaggcagac 1320  
taccactttag agtatgagga actgtttgaa aacctggcag agattgaaa ggcgaagat 1380  
atgaatgaaa ctagaacaac tcttcttgat cctgattcct gcagaaaggc cataactgtg 1440  
tatagtgaa gaggacactt cgtggattca tgggtgactc cgggctttga ctccaggag 1500  
caatgcaccc agaaggctgc cgaaggatat acctagttct actatgtgga tgtcttgat 1560  
gggaagctgg cctgtgtgaa caagtgcacc aaaggaaaga agtcgcaaat gaactgtaac 1620  
ctgggcacat gtcatcgtca acgcagtga ccccgctgc ctgtgcccac atacgaacac 1680  
acactgggtac tggggagaga cctgtgaatt caacatcgcc aagagcctcg tgtatggat 1740  
cgtgggggct gtgatggcgg tgcgtgtgtc cgcattgac atcctaatac tcttatcag 1800  
cctatcccg agaaaacggc acagggaaca gtatgatgt cctcaagagt ggcgaaggga 1860  
aggcaccctt ggcatcttcc agaagacggc catctgggaa gaccagaatc tgagggagag 1920  
cagattcggc ctggaagacg cctacacaaa ctcccgccc accctggaga ctgttgactc 1980  
tggcacagag ctccacatcc agaggccgga gatggtaqca tccactgtgt gaggcaacgg 2040  
gggctccca cctcactcct gctcgttca ggagagctgc aacacagag ctccacacaa 2100  
gctcccggg cgggtccaga ggagacggaa gtccagccct gaagccgtgc ctgtctgag 2160  
ctgacagact tggccagtc cctgctgtg ctccctgtgg ggaagcctgg gggctgtgag 2220  
cctctccatc cgggagcttc cagactccca gaagcctcg cactccgtc tctccctggg 2280  
tggctcccca ctctggaaat tccctaccaa taaaagcaaa tctgaaagct c 2331

&lt;210&gt; 9

&lt;211&gt; 909

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 9

gaggagggtg ggcgaacag acaggcgatt aatgcggctc ttaccagggc aaccaggact 60  
acagtataca ttgtggacat tcaggacata gattctgcag ctccggcccq acctcactcc 120  
tacctcgatg cctactttgt ctccccaat gggctcagccc tgacccttga tgaactgagt 180  
gtgatgatcc ggaatgata ggaactcgtg acgcagctgc tgcagctggg gctgtgtgtg 240  
ctgggctccc aggagagcca ggagtcagac ctgtcgaaac agctcatcag tgcctcata 300  
ggattgggag tggctttgct gctgttctt gtgatcatga ccattggcct cgtgtgtgtg 360  
cgggaagagct acaaccggaa gcttcaagct atgaaggctg ccaaggaggc caggaaagaca 420  
gcagcagggg tgatgcttct agccctgct atcccaggga ctaactgtga cctctctccc 540  
cgagccaacc ccattgctga cctccccaac aaagacctgg gcttggagta cctctctccc 600  
tccaatgacc tggactctgt cagcgtcaac tccctggacg acaactctgt ggaatgtggac 660  
aagaacagtc aggaactcaa ggagcacagg ccactacaca caccactaga gccagatcca 720  
gagccctga gctgtgtcct gttaggacgg caggcaggcg caagtggaca gctggagggg 780  
ccatccatca ccaacgctgg cctggacacc acggacctgt gctagggggc cccactcttc 840  
tggacccctt gaagaggccc taccacaccc taactgcacc tgttccctg gagatgaaa 900  
tatatgacgc tggcctgctt cctgcttttg gccaatcacg gcagacaggg gctggggaaa 960  
tattttatt 999

&lt;210&gt; 10

&lt;211&gt; 510

&lt;212&gt; PRT

WO 99/60161

PCT/US99/10498

&lt;213&gt; Homo sapiens

&lt;400&gt; 10

Met Arg Pro Gly Leu Ser Phe Leu Leu Ala Leu Leu Phe Phe Leu Gly  
 1 5 10 15

Gln Ala Ala Gly Asp Leu Gly Asp Val Gly Pro Pro Ile Pro Ser Pro  
 20 25 30

Gly Phe Ser Pro Phe Pro Gly Val Asp Ser Ser Ser Ser Phe Ser Ser  
 35 40 45

Ser Ser Arg Ser Gly Ser Ser Ser Ser Arg Ser Leu Gly Ser Gly Gly  
 50 55 60

Ser Val Ser Gln Leu Phe Ser Asn Phe Thr Gly Ser Val Asp Asp Arg  
 65 70 75 80

Gly Thr Cys Gln Cys Ser Val Ser Leu Pro Asp Thr Thr Phe Pro Val  
 85 90 95

Asp Arg Val Glu Arg Leu Glu Phe Thr Ala His Val Leu Ser Gln Lys  
 100 105 110

Phe Glu Lys Glu Leu Ser Lys Val Arg Glu Tyr Val Gln Leu Ile Ser  
 115 120 125

Val Tyr Glu Lys Lys Leu Leu Asn Leu Thr Val Arg Ile Asp Ile Met  
 130 135 140

Glu Lys Asp Thr Ile Ser Tyr Thr Glu Leu Asp Phe Glu Leu Ile Lys  
 145 150 155 160

Val Glu Val Lys Glu Met Glu Lys Leu Val Ile Gln Leu Lys Glu Ser  
 165 170 175

Phe Gly Gly Ser Ser Glu Ile Val Asp Gln Leu Glu Val Glu Ile Arg  
 180 185 190

Asn Met Thr Leu Leu Val Glu Lys Leu Glu Thr Leu Asp Lys Asn Asn  
 195 200 205

Val Leu Ala Ile Arg Arg Glu Ile Val Ala Leu Lys Thr Lys Leu Lys  
 210 215 220

Glu Cys Glu Ala Ser Lys Asp Gln Asn Thr Pro Val Val His Pro Pro  
 225 230 235 240

WO 99/60161

PCT/US99/10498

Pro Thr Pro Gly Ser Cys Gly His Gly Gly Val Val Asn Ile Ser Lys  
245 250 255

Pro Ser Val Val Gln Leu Asn Trp Arg Gly Phe Ser Tyr Leu Tyr Gly  
260 265 270

Ala Trp Gly Arg Asp Tyr Ser Pro Gln His Pro Asn Lys Gly Leu Tyr  
275 280 283

Trp Val Ala Pro Leu Asn Thr Asp Gly Arg Leu Leu Glu Tyr Tyr Arg  
290 295 300

Leu Tyr Asn Thr Leu Asp Asp Leu Leu Leu Tyr Ile Asn Ala Arg Glu  
305 310 315 320

Leu Arg Ile Thr Tyr Gly Gln Gly Ser Gly Thr Ala Val Tyr Asn Asn  
325 330 335

Asn Met Tyr Val Asn Met Tyr Asn Thr Gly Asn Ile Ala Arg Val Asn  
340 345 350

Leu Thr Thr Asn Thr Ile Ala Val Thr Gln Thr Leu Pro Asn Ala Ala  
355 360 365

Tyr Asn Asn Arg Phe Ser Tyr Ala Asn Val Ala Trp Gln Asp Ile Asp  
370 375 380

Phe Ala Val Asp Glu Asn Gly Leu Trp Val Ile Tyr Ser Thr Glu Ala  
385 390 395 400

Ser Thr Gly Asn Met Val Ile Ser Lys Leu Asn Asp Thr Thr Leu Gln  
405 410 415

Val Leu Asn Thr Trp Tyr Thr Lys Gln Tyr Lys Pro Ser Ala Ser Asn  
420 425 430

Ala Phe Met Val Cys Gly Val Leu Tyr Ala Thr Arg Thr Met Asn Thr  
435 440 445

Arg Thr Glu Glu Ile Phe Tyr Tyr Tyr Asp Thr Asn Thr Gly Lys Glu  
450 455 460

Gly Lys Leu Asp Ile Val Met His Lys Met Gln Glu Lys Val Gln Ser  
465 470 475 480

Ile Asn Tyr Asn Pro Phe Asp Gln Lys Leu Tyr Val Tyr Asn Asp Gly  
485 490 495



WO 99/60161

PCT/US99/10498

Tyr	Leu	Leu	Asn	Tyr	Asp	Leu	Ser	Val	Leu	Gln	Lys	Pro	Gln
			500					505					510

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/10498

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) : C12Q 1/68; G01N 33/53, 33/574

US CL : 435/4, 6, 7.1

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/4, 6, 7.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Seisearch, HCAPLUS, medline, biosis, medbase, wpids, jicst-plus, biobusiness, biotechds, phis, phic, embas

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96/39419 A1 (HUMAN GENOME SCIENCES, INC.) 12 December 1996, see entire document.	1-5

☐ Further documents are listed in the continuation of Box C. ☐ See parent family annex.

- |  |  |
|--|--|
| <ul style="list-style-type: none"><li>* Special categories of cited documents:</li><li>*A* document defining the general state of the art which is not considered to be of particular relevance</li><li>*E* earlier document published on or after the international filing date</li><li>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</li><li>*O* document referring to oral disclosures, use, exhibition or other means</li><li>*P* document published prior to the international filing date but later than the priority date claimed</li></ul> | <ul style="list-style-type: none"><li>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</li><li>*X* documents of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</li><li>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</li><li>*Z* document member of the same patent family</li></ul> |
|--|--|

Date of the actual completion of the international search

16 AUGUST 1999

Date of mailing of the international search report

10 SEP 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

NANCY JOHNSON

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/10498

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 6  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
  
The sequence listing in computer readable form did not comply with required standards. The disc could not be preprocessed. Thus, a search of claim 6, drawn to SEQ ID NO:3, 4, 5 or 7, could not be carried out.
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐  
☐

The additional search fees were accompanied by the applicant's protest.  
No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet(1)XJuly 1992)\*